

The intricate relationship between gut and kidney

A intrincada relação entre o intestino e os rins

Authors

Regiane S. Cunha¹
Andréa E. Stinghen¹

¹ Universidade Federal do Paraná,
Departamento de Patologia
Básica, Laboratório de Nefrologia
Experimental, Curitiba, PR, Brasil.

In chronic kidney disease (CKD), the intestinal epithelium has compromised functions, including its ability to act as a barrier and to contribute to gut homeostasis. Data from the literature have shown that changes in the gut barrier contribute to systemic inflammation in patients with CKD. However, these pathological processes have not yet been fully elucidated. The dysfunction of the intestinal epithelium can be explained, at least in part, by the effects induced by uremia resulting from CKD on intestinal epithelial cells, such as enterocytes and colonocytes¹⁻⁷.

Considering the crucial role of the intestinal epithelium barrier as well as its intrinsic relationship with the immune system, Andrade *et al.* evaluated the effects of uremic serum on colonocytes in vitro, a study published in the present issue of the Brazilian Journal of Nephrology. For this purpose, the authors incubated colonocytes with the serum of healthy individuals, patients on conservative treatment, and patients on hemodialysis (HD), pre- and post-HD. One of the main findings of this study is the significant increase of proinflammatory cytokine interleukin-6 (IL-6) expression by the colonocytes that were incubated with the serum of both pre- and post-HD patients. Corroborating these findings, Lau *et al.* found an increase in the expression of inflammatory molecules, such as the monocyte chemoattractant protein-1 (MCP-1) and cyclooxygenase-2 (COX-2) proteins, and a greater infiltration of leukocytes in the colon of nephrectomized rats¹. These data, therefore, demonstrate that the uremic environment is able to modulate the inflammatory response in the gut barrier. In addition, Andrade *et al.* did not observe a significant difference in the production of reactive oxygen species (ROS) and expression of toll-like receptors (TLR), proteins that recognize bacterial components and are important for gut homeostasis⁷.

The intestinal epithelium may present an impairment of the intercellular junctions and consequent increase in permeability under pathological conditions. Andrade *et al.* investigated this process through the evaluation of transepithelial electrical resistance of the colonocytes, but no significant difference was found when the cells were exposed to uremic serum. However, Vaziri *et al.* demonstrated that colonocytes exposed to uremic serum from HD patients presented low transepithelial resistance and a significant reduction in the expression of intercellular junction proteins, such as claudin-1, occludin, and zonula occluden-1 (ZO-1)². In another study conducted by Vaziri *et al.*, the same effect was observed when colonocytes were exposed to urea, a uremic toxin that has been extensively studied in recent years³. Similarly, Lau *et al.* observed in vivo a reduction of claudin-2 expression in colonic tissue of nephrectomized rats⁴. Together, these studies strongly suggest that uremia affects negatively the integrity of the gut barrier.

Gut barrier dysfunction allows bacterial translocation, which is the entry of bacteria and products derived from the microbiota, such as endotoxins, to the bloodstream. In fact, clinical studies have demonstrated increased bacterial translocation that is also associated with inflammation in patients with CKD⁴. Moreover, endotoxins, in particular, are associated with hypervolemia in patients with CKD and experimental data have also shown that endotoxins can lead to the activation of the immune system^{5,6}.

Regarding intestinal homeostasis, the gut microbiota should be considered since it is associated with epithelial cells and is responsible for the formation of various uremic toxins and their precursors. Exemplifying this multifaceted panorama, we can cite the uremic toxins *p*-cresyl sulfate and indoxyl sulfate, closely associated with inflammation and cardiovascular diseases

Submitted on: 02/23/2018.

Approved on: 04/22/2018.

Correspondence to:

Andréa E. Stinghen.
E-mail: andreastinghen@ufpr.br

DOI: 10.1590/1678-4685-JBN-2018-00020001



in patients with CKD. In addition, clinical studies have shown that the composition of the microbiota is altered in patients with CKD, with a greater abundance of groups capable of producing uremic toxins. Thus, in recent years, several studies with therapeutic focus have been developed in order to modulate the gut microbiota of patients through prebiotics, postbiotics, and symbiotics, but are still inconclusive and further investigation is necessary^{4,7}.

Based on the study conducted by Andrade *et al.* and in the literature, it seems reasonable to suggest that uremia affects the intestinal epithelium by inducing the inflammatory process and compromising the intercellular junctions.⁸ However, little is known about intestinal epithelial dysfunction and CKD, and recently the research group of Andrade has published a review article highlighting the need to better understand the relationship between intestinal epithelium, microbiota, and CKD⁷. Therefore, further studies are needed to elucidate the complex cellular and molecular mechanisms modulated by uremia in the intestinal epithelium. Finally, findings in this area may contribute to the development of new therapeutic targets in order to improve patient survival.⁸

REFERENCES

1. Lau WL, Liu SM, Pahlevan S, Yuan J, Khazaeli M, Ni Z, et al. Role of Nrf2 dysfunction in uremia-associated intestinal inflammation and epithelial barrier disruption. *Dig Dis Sci* 2015;60:1215-22. DOI: 10.1007/s10620-014-3428-4
2. Vaziri ND, Goshtasbi N, Yuan J, Jellbauer S, Moradi H, Raffatellu M, et al. Uremic plasma impairs barrier function and depletes the tight junction protein constituents of intestinal epithelium. *Am J Nephrol* 2012;36:438-43. DOI: 10.1159/000343886
3. Vaziri ND, Yuan J, Norris K. Role of urea in intestinal barrier dysfunction and disruption of epithelial tight junction in chronic kidney disease. *Am J Nephrol* 2013;37:1-6. DOI: 10.1159/000345969
4. Wang F, Jiang H, Shi K, Ren Y, Zhang P, Cheng S. Gut bacterial translocation is associated with microinflammation in end-stage renal disease patients. *Nephrology (Carlton)* 2012;17:733-8. DOI: 10.1111/j.1440-1797.2012.01647.x
5. Gonçalves S, Pecoits-Filho R, Perreto S, Barberato SH, Stinghen AE, Lima EG, et al. Associations between renal function, volume status and endotoxaemia in chronic kidney disease patients. *Nephrol Dial Transplant* 2006;21:2788-94. DOI: 10.1093/ndt/gfl273
6. Hauser AB, Stinghen AE, Gonçalves SM, Bucharles S, Pecoits-Filho R. A gut feeling on endotoxemia: causes and consequences in chronic kidney disease. *Nephron Clin Pract* 2011;118:c165-72. DOI: 10.1159/000321438
7. Andrade LS, Ramos CI, Cuppari L. The cross-talk between the kidney and the gut: implications for chronic kidney disease. *Nutrire* 2017;42:27. DOI: 10.1186/s41110-017-0054-x
8. Andrade LS, Dalboni MA, Carvalho JTG, Grabulosa CC, Pereira NBF, Aoike DT, Cuppari L. In vitro effect of uremic serum on barrier function and inflammation in human colonocytes. *J Bras Nefrol* 2018;40:2016-23.

ERRATUM

In the article “*The intricate relationship between gut and kidney*” with DOI code number <http://dx.doi.org/10.1590/1678-4685-JBN-2018-00020001> published at Brazilian Journal of Nephrology in 2018:

Where it was written:

Based on the study conducted by Andrade et al. and in the literature, it seems reasonable to suggest that uremia affects the intestinal epithelium by inducing the inflammatory process and compromising the intercellular junctions. However (...)

Should read:

Based on the study conducted by Andrade et al. and in the literature, it seems reasonable to suggest that uremia affects the intestinal epithelium by inducing the inflammatory process and compromising the intercellular junctions.⁸ However (...)

Where it was written:

7. Andrade LS, Ramos CI, Cuppari L. The cross-talk between the kidney and the gut: implications for chronic kidney disease. *Nutrire* 2017;42:27. DOI: 10.1186/s41110-017-0054-x

Should read:

7. Andrade LS, Ramos CI, Cuppari L. The cross-talk between the kidney and the gut: implications for chronic kidney disease. *Nutrire* 2017;42:27. DOI: 10.1186/s41110-017-0054-x
8. Andrade LS, Dalboni MA, Carvalho JTG, Grabulosa CC, Pereira NBF, Aoike DT, Cuppari L. In vitro effect of uremic serum on barrier function and inflammation in human colonocytes. *J Bras Nefrol* 2018;40:2016-23.